

References

1. Balunas MJ, Su B, Riswan S, Fong HH, Brueggemeier RW, Pezzuto JM, Kinghorn AD. Isolation and Characterization of Aromatase Inhibitors from *Brassaiopsis glomerulata* (Araliaceae). *Phytochem Lett*. 2009 Feb 19;2(1):29-33.
2. Jung BI, Kim MS, Kim HA, Kim D, Yang J, Her S, Song YS. Caffeic acid phenethyl ester, a component of beehive propolis, is a novel selective estrogen receptor modulator. *Phytother Res*. 2010 Feb;24(2):295-300.
3. Collins NH, Lessey EC, DuSell CD, McDonnell DP, Fowler L, Palomino WA, Illera MJ, Yu X, Mo B, Houwing AM, Lessey BA. Characterization of antiestrogenic activity of the Chinese herb, *prunella vulgaris*, using in vitro and in vivo (Mouse Xenograft) models. *Biol Reprod*. 2009 Feb;80(2):375-83.
4. Wormke M, Stoner M, Saville B, Walker K, Abdelrahim M, Burghardt R, Safe S. The aryl hydrocarbon receptor mediates degradation of estrogen receptor alpha through activation of proteasomes. *Mol Cell Biol*. 2003 Mar;23(6):1843-55.
5. Klinge CM, Kaur K, Swanson HI. The aryl hydrocarbon receptor interacts with estrogen receptor alpha and orphan receptors COUP-TFI and ERalpha1. *Arch Biochem Biophys*. 2000 Jan 1;373(1):163-74.
6. Kempaiah RK, Srinivasan K. Influence of dietary curcumin, capsaicin and garlic on the antioxidant status of red blood cells and the liver in high-fat-fed rats. *Ann Nutr Metab*. 2004 Sep-Oct;48(5):314-20.
7. Masuda T, Hidaka K, Shinohara A, Maekawa T, Takeda Y, Yamaguchi H. Chemical studies on antioxidant mechanism of curcuminoid: analysis of radical reaction products from curcumin. *J Agric Food Chem*. 1999 Jan;47(1):71-7.
8. Bhutani MK, Bishnoi M, Kulkarni SK. Anti-depressant like effect of curcumin and its combination with piperine in unpredictable chronic stress-induced behavioral, biochemical and neurochemical changes. *Pharmacol Biochem Behav*. 2009 Mar;92(1):39-43.
9. Scapagnini G, Foresti R, Calabrese V, Giuffrida Stella AM, Green CJ, Motterlini R. Caffeic acid phenethyl ester and curcumin: a novel class of heme oxygenase-1 inducers. *Mol Pharmacol*. 2002 Mar;61(3):554-61.
10. Balogun E, Hoque M, Gong P, Killeen E, Green CJ, Foresti R, Alam J, Motterlini R. Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive element. *Biochem J*. 2003 May 1;371(Pt 3):887-95.
11. Jeong GS, An RB, Pae HO, Oh GS, Chung HT, Kim YC. Heme oxygenase-1 inducing constituent of *Prunella vulgaris* in HepG2 cells. *Biol Pharm Bull*. 2008 Mar;31(3):531-3.
12. Stone M, Ibarra A, Roller M, Zangara A, Stevenson E. A pilot investigation into the effect of maca supplementation on physical activity and sexual desire in sportsmen. *J Ethnopharmacol*. 2009 Dec 10;126(3):574-6.
13. Suresh D, Srinivasan K. Influence of curcumin, capsaicin, and piperine on the rat liver drug-metabolizing enzyme system in vivo and in vitro. *Can J Physiol Pharmacol*. 2006 Dec;84(12):1259-65.

